


į

MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS - 1963 - A

AD A139396






A Response Model for Multiple Choice Items

David Thissen and Lynne Steinberg University of Kansas

Psychometric Technical Report No. 1





B



Approved for public releases Distribution Unlimited

OOG

84 03 23 006

IC FILE COPY

TORRESSENT TORRESSENT

UNIVERSITY OF CHICAGO

A Response Model for Multiple Choice Items

David Thissen and Lynne Steinberg University of Kansas

Psychometric Technical Report No. 1



This research was sponsored by the Personnel and Training Research Program, Psychological Sciences Division, Office of Naval Research under Contract No. N00014-83-C-0283, Contract Authority Identification No. NR 475-018.

Approved for public release; distribution unlimited. Reproduction in whole or part is permitted for any purpose of the United States Government.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM
[-,, · · · · · · · · · · · · · · · · ·	3. RECIPIENT'S CATALOG NUMBER
Psychometric Technical Report No. 1 AD-A139396	
4. TITLE (and Subtitle)  //EMS	5. TYPE OF REPORT & PERIOD COVERED
A Response Model for Multiple Choice <del>Models</del>	
	6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(a)	S. CONTRACT OR GRANT NUMBER(s)
David Thissen and Lynne Steinberg	1100014-83-C-0283
· ·	
9. PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
NORC	62763N RF63521803
6030 S. Ellis Avenue	RF63521 NR 475-018
Chicago IL 60637 11. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE
	December 1983
Personnel and Training Research Programs	13. NUMBER OF PAGES
Office of Naval Research (Code 442PT) Arlington, VA. 22217	43
14. MONITORING AGENCY NAME & ADDRESS(II dillerent from Controlling Office)	15. SECURITY CLASS. (of this report)
	Unclassified
	15. DECLASSIFICATION DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)	
Approved for public release; distribution unlimited	
Approved for public release, discribation unimitted	
12 DAG COLUMN CO	
17. DISTRIBUTION STATEMENT (of the abetract entered in Block 20, if different from Report)	
18. SUPPLEMENTARY HOTES	
·	
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)	
	į
20. ABSTRACT (Continue on reverse elde if nece-very and identify by block number)	
An extension of the Bock-Samejima model for multiple choice items is	
introduced. The model provides for varying proba	
alternative when the examinee guesses. A marginal maximum likelihood method	
is devised for estimating the item parameters, and likelihood ratio tests for comparing more and less constrained forms of the model are provided. Appli-	
cations of the model are illustrated with item response data for the word	
knowledge and general science subtests of the Armed Services Vocational	
Aptitude Battery	

DD 1 JAN 73 1473


A RESPONSE MODEL FOR MULTIPLE CHOICE ITEMS

David Thissen and Lynne Steinberg

University of Kansas

#### **ABSTRACT**

An extension of the Bock-Samejima model for multiple choice items is introduced. The model provides for varying probabilities of the response alternative when the examinee guesses. A marginal maximum likelihood method is devised for estimating the item parameters, and likelihood ratio tests for comparing more and less constrained forms of the model are provided. Applications of the model are illustrated with item response data for the word knowledge and general science subtests of the Armed Services Vocational Aptitude Battery.



# A Response Model for Multiple Choice Items

In practical ability measurement, binary item response models have been applied routinely to data from multiple-choice tests with four or five alternatives per item. To use the binary models, the data have been dichotomized (correct and incorrect) and the distinct identity of the incorrect alternatives has been lost. This procedure logically follows the tradition of "scoring" a test by considering only the correct answers; however, the binary models are incomplete as theories of the item responses as useful information may be lost in the dichotomization.

It is possible to conceive of a unidimensional latent variable model which completely explains the item response data, and several have been proposed. The model which will be introduced here is an extension of that proposed by Bock (1972) and subsequently extended by Samejima (1979); a different parameterization of the same model has been discussed by Sympson (1983). Bock (1972), Thissen (1976), Sympson (1983), and Levine and Drasgow (1983) have shown that some increased precision of measurement may be obtained when information in the incorrect alternatives on multiple choice tests is included in the item response model. In a subsequent section, we will discuss this gain in information with a new model, as well as some attendant difficulties. In the next section we will introduce the model and describe a workable scheme for maximum likelihood estimation of its parameters.

## The Model

The model for multiple-choice items to be described makes heavy use of the multivariate logistic transformation suggested as an item response model by Bock (1972), so it is useful to begin with a clear understanding of that model. For m categorical responses, we specify m response functions  $z_k = a_k\theta + c_k$ , each of which is a linear function of the latent ability variable  $\theta$ . Usually  $z_{correct}$  would have a positive slope and the other  $z_k$ 's would have lesser positive or negative slopes. The linear functions describe one of the simplest possible relationships between the response and  $\theta$ . To make this a model for categorical item responses, the  $z_k$  (which lie on the real line) must be mapped into [0,1]. This is accomplished by the multivariate logistic transformation, so

$$P(x_{j}=h|\theta;\underline{a},\underline{c}) = \frac{\exp(z_{h})}{\sum \exp(z_{k})} . (1)$$

$$k=1,m_{j}$$

The function (1) is Bock's (1972) model for an item response,  $x_j=h$ , in which  $h=1,2,\ldots m_j$  for a multiple choice item j with  $m_j$  (classes of) response alternatives. The item parameters are the vectors  $\underline{a}$  and  $\underline{c}$ , subject to two suitable linear constraints (see below), giving  $2(m_j-1)$  free parameters. The model described by (1) is moderately flexible, but lacks flexibility in certain crucial respects at the extremes. Specifically, one of the response alternatives must have the largest positive value of  $a_k$ ; the trace line for that alternative is then monotonic increasing. That may be theoretically acceptable, since it

is probably the correct response. However, another alternative must have the largest negative value of  $a_k$  ( $\Sigma a_k$ =0 is one of the linear constraints), and that alternative's trace line must be monotonic decreasing. The latter aspect of the model is less acceptable: it implies that as ability decreases the probability of selecting one particular incorrect response approaches unity, and all the others go to zero. That is unlikely.

Samejima (1979) proposed a solution to this problem in a conceptual modification of Bock's (1972) model in which she added a completely latent response category labelled "zero." Here, we will sometimes refer to this category as "don't know" (DK). Lord (1982) has introduced a similar conceptual entity which he describes by saying the examinee is "(totally) undecided." In Samejima's model, DK is not an observed response, but a latent one, multi-logit-linearly dependent on a latent trait (giving two layers latent); the idea was that some proportion d<sub>h</sub> (=1/m<sub>j</sub> in Samejima (1979)) "guessed" each of the observable response alternatives and were "mixed in" with the examinees who chose those alternatives intentionally. So the model becomes

$$P(x_j = h \mid \theta; \underline{a}, \underline{c}, \underline{d}) = \frac{\exp(z_h)}{\sum \exp(z_k)} + \frac{d_h \exp(z_0)}{\sum \exp(z_k)}$$

$$k = 0, m_j + \frac{\sum \exp(z_k)}{k = 0, m_j}$$

in which the second term adds a proportion  $d_h$  of those who "don't know" into each trace line; call it  $P_i(h)$  for future brevity:

$$P_{j}(h) = \frac{\exp(z_{h}) + d_{h} \exp(z_{0})}{\sum_{k=0, m_{j}} \exp(z_{k})},$$
 (2)

in which h takes the values  $1,2,\ldots m_j$ . Samejima's (1979) model had two more parameters than Bock's (1972) model:  $a_0$  and  $c_0$ . In her presentation, the  $d_h$  were fixed and equalled  $1/m_j$ ; this represents the hypothesis that those of sufficiently low ability assign their responses randomly with equal probability to each of the response alternatives. We found that unlikely; later we will show that it is not empirically the case. So we extend the version of the model given in (2) to allow the  $d_h$ ,  $h=1,2,\ldots m_j$  to be functions of estimated parameters.

## Indeterminacies and Constraints

The model expressed in (2) has a number of indeterminacies and requires the imposition of some constraints to become identifiable. The linear constraints required on a and c,

$$\Sigma_{a} = \Sigma_{c} = 0$$

$$k=0, m_{j}^{k} = 0, m_{j}^{k}$$

are imposed by reparameterization:

$$a = T\alpha$$

and

$$c = T\gamma$$
,

where T is an  $(m_j+1)$  x  $m_j$  transformation matrix, and  $\underline{\alpha}$  and  $\underline{\gamma}$  are vectors of free parameters of order  $m_j$ ; T is the transpose of that in Bock (1972) throughout. There is a further indeterminacy with respect to the sign of  $\underline{\alpha}$ ; this is an indeterminacy of reflection of the latent variable

 $\theta$  which is logically identical to the rotational indeterminacy of common factor analysis. The reflection indeterminacy is not explicitly solved, but rather the iterative solution of the likelihood equations is started with  $a_h$  positive for h = the correct response; this is usually sufficient to keep the estimated solution "right-side-up."

Since the vector  $\underline{\mathbf{d}}$  represents a set of  $\mathbf{m}_j$  proportions, the  $\mathbf{d}_h$  must be constrained to sum to unity and lie on the interval [0,1]. The dual constraint is imposed in two parts. In the first, the same multivariate logistic used in the model is also employed to make the  $\mathbf{d}_h$  proportions from a set of psuedo-parameters  $\mathbf{d}_h$ , which lie on the real line:

$$d_{\mathbf{h}} = \exp(d_{\mathbf{h}}^{*}) / \sum_{k=1, m, j} \exp(d_{\mathbf{k}}^{*}) . \qquad (3)$$

The parameters  $d_h^*$  are on the real line, but like  $\underline{a}$  and  $\underline{c}$ , they must satisfy  $\Sigma d_k^* = 0$ . So

$$\underline{\mathbf{d}}^* = \mathbf{T}_2 \underline{\delta} \tag{4}$$

in which  $T_2$  is an  $m_j \times (m_j-1)$  transformation matrix of the same form as T (above) but of lesser order, and  $\underline{\delta}$  is a vector of length  $(m_j-1)$  of parameters which (through (4) and (3)) give the proportions  $d_h$ .

So the set of free parameters for an item consists of m $_j$   $\alpha_k$ 's ("a-contrasts"), m $_j$   $\gamma_k$ 's ("c-contrasts"), and  $(m_j-1)$   $\delta_k$ 's ("d-contrasts"), for a total of  $3m_j-1$  parameters: 11 for a fouralternative multiple-choice item.

The estimation system we use permits any of the elements of  $\{\underline{\alpha},\underline{\gamma},\underline{\delta}\}$  to be fixed at any constant value, or to be constrained to be

equal to any other parameter. With this facility, our model (2) includes the previous models hierarchically as subsets. If  $\underline{\delta}$  is fixed at  $\underline{0}$ , than all the  $d_h=1/m_j$  and (2) becomes Samejima's (1979) model. If the parameters  $a_0$  and  $c_0$  are fixed at zero, the  $d_h$  become irrelevant (and are deleted mechanically) and the model becomes Bock's (1972) model. In our examples, we will compare the goodness of fit of all of these models, as well as versions of (2) that include equality constraints on particular parameters imposed across items.

## The Relationship of the Model to Data

Models in item response theory (IRT) are intended to explain observed covariation among test-item responses. Unidimensional "latent trait" theories explain that covariation by appeal to an underlying latent variable (usually denoted  $\theta$ ) on which the probabilities of the responses are functionally dependent. Each pair of item responses is theorized to have non-zero covariance because the probability of both item responses depends on  $\theta$ . The models usually hold that  $\theta$  is the sole cause of covariation among the item responses; conditional on  $\theta$ , the item responses are theorized to be independent ("local independence").

A complete IRT model for the item response data for a multiple-choice test in which n items each have m alternatives would be a model for the covariation in the  $m^0$  contingency table containing the counts of respondents giving each response pattern. Such a  $4^2$  table for two

four-alternative multiple-choice items is given in Table 1.

44444

## Insert Table 1 About Here

Evidence that there is some information in the examinees' i rrect choices is apparent (or at least can be found) in the covar ace in the table. (The column percentages for the counts are given i prentheses to aid in interpretation.) Alternative C is correct for item 2; respondents who select C on item 2 are most likely to choose B on item 1; B is correct, so that is double evidence of their high ability. But examinees who choose A or B on item 2 are more likely to respond correctly to item 1 (36% and 34%) than those who pick D on item 2 (only 25%). A possible explanation is that those who select A or B on item 2 are higher on the ability continuum than are those who pick D. Even with 16 cells, this sort of argument may be lengthy, but the idea is that wrong responses are related to each other; if the responses are locally independent, this implies a relationship to "the trait."

The goodness of fit of a multiple category IRT model to the data in such a table may be evaluated with conventional likelihood ratio  $\chi^2$  statistics. For sets of data with more examinees than cells in the table, the likelihood ratio test against a general multinomial alternative may be used. For larger tables (35 four-choice items gives a  $4^{35}$  table), there is no "general multinomial alternative"; but the likelihood ratio test between hierarchically nested models still may be used to evaluate the significance of the additional parameters of the

larger model.

### Estimation

In general, the data for the estimation of the parameters  $\{\underline{\alpha},\underline{\gamma},\underline{\delta}\}_j$  for each item j, j=1,...,n consist of the counts of response patterns  $\mathbf{r}_{\underline{\mathbf{x}}}$  (where each  $\mathbf{x}_j$  may take values in  $[1,2,\ldots m_j]$ ) in an  $\mathbf{m}_1\times\mathbf{m}_2\times\ldots\times\mathbf{m}_n$  contingency table like Table 1. The probability of observing a particular response pattern  $\underline{\mathbf{x}}$  when drawing an examinee from a population in which  $\theta$  follows a distribution  $\phi(\theta)$ , assumed N(0,1) in the examples, is

$$P(\underline{x}) = \int_{-\infty}^{\infty} \prod_{j=1,n} P_j(x_j) \phi(\theta) d\theta$$
 (5)

in which  $P_{j}(x_{j})$  is from equation (2).

The likelihood for the entire set of observed data is

$$L = C \prod_{\underline{x}} P(\underline{x})^{r} \underline{x}$$
 (6)

in which C is a normalizing constant not dependent on the parameters, and the product runs over all possible response patterns. In practice, of course, for more than a handful of items the number of possible response patterns is astronomical, and the "count" in each cell is one or zero. In such cases, only those response patterns actually observed need to be considered in the computation of (6), or (7), a function proportional to the loglikelihood:

$$\ell_0 \sim \frac{\Sigma}{\underline{x}} r_{\underline{x}} \log P(\underline{x})$$
 (7)

For a very small number of items (3 or 4; the parameters of the full model are not identifiable with only 2 items, as there are fewer cells than parameters), it may be possible to obtain the MLEs by directly maximizing (7) with a Newton-type algorithm. Bock (1972) reported such results for his original model with procedures similar to those used by Bock and Lieberman (1970).

Bock and Aitkin (1981) described an extremely simple and elegant algorithm for maximizing functions like (7) for binary models, and we extend that algorithm to the estimation of multiple category models. The Bock-Aitkin algorithm is a two-step procedure like the "EM-algorithm" (Dempster, Laird, and Rubin, 1977), so we describe first the "E-step" and then the "M-step." The procedure is iterative, and EM-pairs are repeated until the process converges.

## The E-step

The Bock-Aitkin algorithm is based on a discrete representation of  $\phi(\theta)$  and the integrand of  $P(\underline{x})$ , both continuous densities, over Q "quadrature points"  $\theta_q$ , with q=1,2,...Q. Such a discrete representation of the continuous densities may be made arbitrarily close to continuous reality by choosing Q large, just as numerical integration may be made arbitrarily accurate by using sufficient quadrature points. However, large values of Q slow the computations; Q=10 over the range  $\theta$ =-4.5 to  $\theta$ =4.5 in unit steps seems to be sufficiently accurate and fast for many applications and is used in all of the examples here.

Under the assumption that the population is composed of individuals who are members of Q discrete "classes" with values  $\theta_1$ ,  $\theta_2$ , ...,  $\theta_Q$  on the latent variable, "complete data sufficient statistics" for the estimation of the item parameters  $\{\underline{\alpha},\underline{\gamma},\underline{\delta}\}_j$  for item j would consist of a table of counts  $R_j^+$ , in which each element  $r_{jkq}^+$  is the number of individuals in class  $\theta_q$  selecting response alternative k on item j. So the E-step of the Bock-Aitkin algorithm consists of computing the expected values of the  $r_{jkq}^+$ , conditioned on the data and the current provisional estimates of the item parameters.

Using provisional estimates of the item parameters for each item (as starting values we use  $\underline{\delta}=\underline{0}$ ,  $\underline{\gamma}=\underline{0}$ , and  $\alpha_k=1$  for categories k which are incorrect and  $\alpha_k=2$  for category k correct), compute the elements of the  $m_i$  x Q table  $R_i^*$  containing:

$$E(r_{jkq}^{*} | data; \{\underline{\hat{\alpha}}, \underline{\hat{\gamma}}, \underline{\hat{\delta}}\}) = \sum_{(x \text{ in } \xi)} r_{\underline{x}} [P(\underline{x}; \theta_{q}) / \sum_{q} P(\underline{x}; \theta_{q})]$$
(8)

in which  $\xi$  is the set of  $\underline{x}$  in which  $x_{\,\underline{i}} \! = \! k$  and

$$P(\underline{x};\theta_{q}) = \prod_{j=1,n} P_{j}(x_{j})\phi(\theta_{q}) . \qquad (9)$$

Note that, while (9) and (8) are computed in a (potentially long) loop over the observed response patterns, the values  $P_j(k)$  for each item from (2) are required only for a fixed set of Q values of  $\theta$ . If those values are placed in a table before the E-step is begun, the computations involved in (8) and (9) are limited to table look-up, multiplication, and addition. The E-step yields a set of n m<sub>j</sub> x q tables of non-integral artificial "counts" which are used as data in the M-step.

## The M-step

The M-step consists of maximum likelihood estimation of the parameters  $\{\underline{\alpha},\underline{\gamma},\underline{\delta}\}_j$  for all items j=1,2,...n, using the tables of expected values  $R_j^*$  as data. It is simply nonlinear regression.

in which  $P_j(k;\theta_q)$  is equation (2) evaluated at  $\theta_q$ . Standard gradient methods may be used to maximize  $\ell_j$  over the parameter space. The MLEs of the parameters are obtained where

$$\frac{\partial \ell_j}{\partial \zeta} = 0$$

for all parameters in  $\zeta = \{\underline{\alpha}, \underline{\gamma}, \underline{\delta}\}$ . Since for any set of parameters constrained by T such that  $\kappa = T\zeta$ ,

$$\frac{\partial \ell_{j}}{\partial \zeta} = T \frac{\partial \ell_{j}}{\partial \kappa} , \qquad (11)$$

we will state the required components of the gradient for the constrained parameters  $\{\underline{a},\underline{c},\underline{d}\}$ ; the gradients used to maximize (10) are then given by (11). Further,

$$\frac{\partial \ell_{j}}{\partial \kappa} = k q^{\sum_{j} \sum_{k \neq j} r_{jkq}^{k}} \frac{1}{P_{j}(k;\theta_{q})} \frac{\partial P_{j}(k;\theta_{q})}{\partial \kappa} ; \qquad (12)$$

So what we really need are the derivatives of (2) with respect to  $\underline{a}$ ,  $\underline{c}$ , and  $\underline{d}^*$  (substitute all for  $\kappa$  above). To write these, a useful bit of

shorthand is

$$e_k = \exp(a_k \theta + c_k)$$
,

which is always assumed to be evaluated at the appropriate value of  $\theta$ ; all summations of  $e_k$  are over categories  $k=0,1,\ldots m_j$ . The elements of the gradient  $(P_h$  is short for  $P_j(h;\theta_q)$  hereafter) to be substituted in (12), then (11), and finally zeroed to maximize (10) are:

$$\frac{\partial P_h}{\partial a_0} = \frac{\left[ (\Sigma e_k) \theta d_h e_0 - (e_h + d_h e_0) \theta e_0 \right]}{(\Sigma e_k)^2}$$

$$\frac{\partial P_h}{\partial a_h} = \frac{\left[ (\Sigma e_k) \theta e_h - (e_h + d_h e_0) \theta e_h \right]}{(\Sigma e_k)^2}$$

$$\frac{\partial P_h}{\partial a_h} = \frac{-\theta e_h}{\sum e_k}$$

$$\frac{\partial P_h}{\partial c_0} = \frac{\left[ (\Sigma e_k) d_h e_0 - (e_h + d_h e_0) e_0 \right]}{(\Sigma e_k)^2}$$

$$\frac{\left(\sum e_{k}\right)e_{h}-\left(e_{h}+d_{h}e_{0}\right)e_{h}}{\left(\sum e_{k}\right)^{2}}$$

$$\frac{\partial P_h}{\partial c_h} = \frac{-e_h}{\sum e_k}$$

$$\frac{\partial P_h}{\partial d_h^*} = \frac{e_1}{\sum e_k} \cdot \frac{\sum d_k^* - d_h^{*2}}{(\sum d_k^*)^2}$$

The gradient does not simplify very much since the model is not part of an exponential family. However, it is possible to locate the maximum of  $\ell_j$  using these derivatives, and use the resulting set of parameters in the next E-step. Our estimation procedure allows equality constraints to be placed across as well as within items; this requires some of the gradients to be accumulated across items, but causes no serious problem beyond book-keeping. We use a conditioned Newton-type algorithm (MINIM, described by Haberman, 1974) to locate the maximum of (10); the conditioning is useful since the matrix of second derivatives may be nearly singular for some items.

## Convergence and Local Minima

Using the updated item parameters from the M-step, the sequence [E-step, M-step] is repeated until either a) the parameters stabilize or b) a fixed number of cycles is reached (we usually use 15 or 20). The algorithm sometimes converges in the parameter space. More often, it does not, but the lack of convergence causes no real problems. In such cases, most of the parameters (and the loglikelihood) remain roughly constant after ten or fifteen cycles, while a parameter (or a few) change linearly and indefinitely. The changing parameter is frequently a particular  $\alpha_k$  (and its associated  $\gamma_k$ ), changing as the associated a rises toward a very high (possibly infinite) MLE. The difference in goodness of fit with such a slope high (3. to 5. or so) and

much higher appears to be negligible, so there is no loss in simply stopping the estimation procedure arbitrarily. In such a case, the trace line is fairly well-determined; but the numerical value of the associated a is not.

A second case in which the parameters fail to clearly converge arises with  $\gamma_k$ 's associated with  $a_k$ 's near zero. The parameter  $c_k$  becomes ill-defined when  $a_k$  is zero, and the associated  $\gamma_k$  "wanders." Again, there is no loss of fit if the estimation procedure is stopped. The Bock-Aitkin algorithm may be speeded slightly using acceleration as in Thissen (1982), but the value of the acceleration parameter should be limited to about -1, which doubles the step-size at each cycle, to avoid oscillation.

The likelihood surface for the model clearly has more then one "local" minimum. The indeterminacy with respect to reflection of  $\theta$  gives two equal, identical modes. With poorly chosen starting values, the estimation procedure has located other (apparent) stationary points; usually they seem to be located in a peculiar region corresponding to some items using both reflections of  $\theta$  in the same solution -- these may be modes or saddle points. Good starting values and the EM-like nature of the Bock-Aitkin algorithm provide a solution in this case, because EM-algorithms only climb local modes. If the algorithm starts near the desired part of the likelihood surface, it will end there. It appears that sensible starting values solve the problem.

The likelihood surface is multi-modal and may be ill-behaved, producing parameter estimates that are sometimes undefined or are on the

boundary. The entire system begs for full Bayesian treatment, with a prior distribution restricting the parameters to a reasonable part of the space under all circumstances. At this stage however, it is not clear what sort of prior may be appropriate. After some experience with the model is gained beyond that in the examples which follow, we may be in a position to specify reasonable parameters for a prior.

Alternatively, a different model, in the exponential family and with a unimodal likelihood (preferably with similar properties to this one in terms of fitting the data), would also solve all of these problems. Such a model has yet to be proposed.

## Characterizing 0

The parameter estimation described above usually has the goal of "calibrating" a set of test items, after which the parameters are to be taken as known and used to characterize  $\theta$  for examinees who produce a particular response pattern  $\underline{\mathbf{x}}$ . Given a set of item parameters, the posterior density for  $\theta$  is

$$P(\underline{x};\theta) = \prod_{j=1,n} P_j(x_j)\phi(\theta) , \qquad (13)$$

in which  $P_j(x_j)$  is from equation (2). If the model is correct, (13) describes the distribution of examinees who respond with pattern  $\underline{x}$ . It can be characterized graphically, and two examples will be presented below in Figure 5.

For more than a few items, (13) is roughly Gaussian in shape, and so it may also be described by estimates of its location and spread. An

extension of the procedure commonly used in binary IRT is to use the mode as an estimate of the location of (13), where

$$\frac{\partial \log P(\underline{x};\theta)}{\partial \theta} = \sum_{j=1,n} \frac{\partial \log P(x_j;\theta)}{\partial \theta} + \frac{\partial \log \phi(\theta)}{\partial \theta}$$

is zero, with the variance approximated by the negative inverse of

$$E = \frac{\partial^2 \log P(\underline{x}; \theta)}{\partial \theta^2}$$

The modal estimate is practical to compute as long as  $\phi(\theta)$  is a reasonable function, and easy if  $\phi(\theta)$  is normal. There is no guarantee that (13) is unimodal; for small numbers of items it is likely to have more than one mode. Multimodality presents potential problems for mechanical use of modal estimates.

However, it is also fairly straightforward to numerically integrate (13) to obtain its mean and variance. The mean has been called the "EAP" (Expected A Posteriori) estimate of  $\theta$  by Bock and Mislevy (1982). An advantage of the EAP procedure over modal estimation is that the derivative of  $\phi(\theta)$  is not required; therefore  $\phi(\theta)$  may take any form describable as a histogram with finite variance.

## Examples

To illustrate application of the model to item response data, we have analyzed several subsets of items from the ASVAB Form 8A administered to a national probability sample of youth by the National Opinion Research Center; the data are described by Bock and Mislevy

(1981). The data used here are from a 10% subsample of the NORC sample, total N=1178. For small illustrations of the performance of the model, we use four-item subsets consisting of the last four items of the Word Knowledge (WK) and General Science (GS) sections of the test. Four items were used because, with roughly a thousand examinees, that is the largest number for which the number of cells in the  $m^n$  table is less than N ( $4^4$ =256), so significance tests of the overall goodness of fit of the model against a general multinomial alternative are feasible. The last items were used because they are most difficult, giving the trace lines more latitude to be fitted.

For the four-item examples, only examinees with complete data (no non-response) for all four items were used. For GS, N=1048 and 214 of the 256 cells of the table were filled (one of the 4 x 4 marginal tables of this table makes up Table 1); for WK, N=976 and only 156 response patterns were observed.

We also fitted several forms of the model to the entire set of 35 WK items, and a subset of 12 of the 25 GS items called "physical science" hereafter. (The ASVAB "General Science" section consists of about half biological and half physical science questions; we selected a subset more likely to be unidimensional.) In these analyses, all examinees were used (N=1178) and non-response data to individual items were ignored (not placed in any category).

#### Results.

Table 2 gives summary goodness of fit results for the four-item examples.

## Insert Table 2 About Here

Bock's (1972) model and Samejima's (1979) modification are both rejected at the p<0.05 level for both sets of data. The model called "ABCD" is a form of (2) in which the vector  $\underline{\delta}$  is estimated, but restricted to be equal across items: this represents the hypothesis that a (constant) proportion of those who "don't know" select alternative A, a different proportion B, and so on; but the distribution over A, B, C, and D (regardless of the "correct" response alternative) is the same for all items. For the WK data the model is a great improvement over Samejima's  $\underline{\delta}=\underline{0}$  model ( $\underline{G}^2=12.6$  on 3 d.f.), and for GS it is not. The estimated vector d for WK for [A,B,C,D] is [.1,.2,.4,.3].

In the last four WK items, D is correct for the first and the last and C is correct for the others; so to test the hypothesis that (somehow) the correct alternative "attracts" guessing we tested "ABCD(C),ABCD(D)" in which the two pairs of items had different vectors  $\underline{\delta}$ . This model fits the WK four-item data, and the improvement over ABCD is significant ( $G^2=10.6$ , d.f.=3, p<0.02); the parameter estimates are given in table 3. The "saturated" version of (2), with 11 free

parameters per item, does not fit the WK data significantly better.

\*

## Insert Tables 3 and 4 About Here

The last four GS items do not have such a convenient structure of correct alternatives; but for these items "correctness" seemed more important than alternative position, so in the model "correct vs. incorrect"  $d_h$  is constrained to be one value (the same for all items) for the correct response and another for all the incorrect responses. This model (barely) fits the GS data. The parameter estimates are given in Table 4. Forty percent of the DK guess the correct alternative; there is no evidence that this varies across items (The last line of Table 2 gives the  $G^2$  for "correct vs. incorrect each item").

When the complete WK data are fitted, the likelihood ratio test against a "general multinomial alternative" for the 1178 observations in a  $4^{35}$  table is, of course, meaningless. Nevertheless, we report likelihood ratio tests between variously constrained versions of the model. The Samejima (1979) model is the most constrained form that we use for comparison. A model of the "ABCD" form of (2), in which  $\underline{\delta}$  is estimated, equal across items, gives a reduction  $G^2$ =41 on 3 d.f., p<0.001; the estimate of  $\underline{d}$  was [.2,.3,.3,.2] for [A,B,C,D]. As with the four-item WK, a model in which  $\underline{\delta}$  is constrained to be equal only among items with the same (letter) correct response fits significantly better:  $G^2$ =72 on 9 d.f., p<0.01. No more complex models were considered, as estimation of 11 parameters per item for the very easy items (to which

the majority of examinees responded correctly) would be estimating many parameters with little data. The estimated parameters for the last four items obtained in the analysis of the entire WK set are given directly below their values for the four-item estimation in Table 3.

Analysis of the physical science portion of the GS section gave similar results: (2) in its "ABCD" form fitted significantly better than Samejima's model,  $G^2=29$  on 3 d.f., p<0.01. The estimate of  $\underline{d}$  was again [.2,.3,.3,.2]. And the model in which  $\underline{d}$  varied according to the letter-value of the correct alternative was better:  $G^2=26$  on 9 d.f., p<0.01. Two of the items in the "physical science" set are among the last four GS items, and their parameter estimates are given in Table 4.

### Discussion

In cases in which the fit can be tested -- with a thousand examinees and four-alternative items, four items -- some forms of the model given by (2) and (5) fit item response data satisfactorily. This represents a major step forward in item analysis, because it is no longer necessary to look at deviations from the fit to examine items; we may examine the fit itself.

Insert Figure 1 About Here

The solid curves in Figure 1 illustrate the trace lines for the "physical science" fit to GS item 23, which is item 2 in Table 1, discussed above. From Table 1 we inferred that those who choose A or B

might be more able than those who chose D. Indeed, the trace line for D restricts that response to those with values of  $\theta<0$  for all practical purposes, while B-responses come mostly from those with  $\theta>0$ , and A is spread all over. The DK trace line indicates that most of those below  $\theta=-.5$  have no idea; and more of them guess correctly than any other way.

The dashed curves in Figure 1 illustrate the trace lines for the "four-item" fit to item 23. Note in Table 4 that the parameter estimates differ a great deal, but the solid and dashed trace lines in Figure 1 differ much less (except for DK) in the middle. DK is a completely inferred latent response and is not really very well-defined by four items. The other curves are more similar in the middle, where the data are; 95% of the population distribution lies between 0s of -2 and +2. The different parameter estimates seem to affect the curves primarily at the extremes.

Unlike algorithms which make use of point-estimates of  $\theta$  in the estimation of item parameters, and require both large numbers of examinees and large numbers of items, the MML estimation procedure used here should be consistent considering the number of examinees alone if the model is correct. That is, as the number of examinees becomes large the estimates for four items or 12 or 35 should all converge to the true values. So the dashed and solid lines in Figure 1 should be the same. They are very similar, and there are three possible reasons for the small differences observed between them. First, one thousand examinees may easily not be "asymptotic" for a table with 256 cells. Second, one or the other solution may not be completely converged; EM-algorithms can

be slow near the maximum. Third, and most likely, the model may be incorrect in that the latent dimension  $\theta$  may be defined slightly differently by the most difficult four items than by the entire test.

Insert Figure 2 About Here

12000 110010 1 110000 11010

Figure 2 similarly illustrates the four-item and complete test estimates of the trace lines for WK 32, which demonstrates this effect more graphically. With estimates from the entire test, DK goes to unity for low  $\theta$ . But the four-item (dashed) curve for DK rises for moderately low  $\theta$  and then goes back down, and A goes to unity for low  $\theta$ . That can't be right; but it is all happening in the region below  $\theta$ =-2, where there are essentially no data and the model is extrapolating. Extrapolation is bad: the model is too flexible at the extremes. Flexibility was one of the goals of the model, but it seems we may have overdone it. This is in marked contrast to traditional IRT models which go to unity, zero, or some asymptote at the extremes, and never misbehave there. Note that even with all of this strangeness, the other trace lines in Figure 2 are essentially identical between  $\theta$ s of -2 and +2.

Insert Figure 3 About Here

Figure 3 shows the trace lines for the correct responses only for WK items 32-35 from the four item set and the whole test, as well as

traditional 3-PL curves estimated for the entire test (dotted lines) for comparison. If the "tail-wagging" to the left of  $\theta$ =-2 is ignored, the pairs of curves for the whole test are nearly identical and the four-item curves are somewhat deviant. That may be capitalization on chance in four items -- or it may be that  $\theta$  differs mildly in the hardest items of the test from its definition in the entire test. None of the curves are extremely different.

## Insert Figure 4 About Here

-----

For WK as a whole, the usual result (of Bock, 1972; Thissen, 1976) is obtained with respect to test information: Figure 4 shows the information curves for the multiple category scoring and the binary scoring (3-PL). For  $\theta$ <0, information from incorrect responses increases total information by about 50%, equivalent to extending the test (for half the examinees) from 35 to 50 items.

## Insert Figure 5 About Here

Figure 5 illustrates the process. The solid curves show the trace lines associated with a particular set of responses [B,B,B,B], all incorrect, for the last four WK items. Then the N(0,1) population distribution is plotted with the the product of all five curves, or posterior density, labelled "Total." The mode of that density is traditionally  $\hat{\theta}$ ; it is about -0.7 (s.e.=0.6) in this case. The dashed

curves show the 3-PL trace lines, all incorrect, so they are all monotonic decreasing. That is not very informative; the (dashed) posterior is broader (s.e.=0.7) and closer to zero ( $\hat{\theta}$ =-0.5) because there is less information so the estimator is "shrunk" more toward the mean of the population distribution. If the model approximates the world reasonably well, the multiple category scoring gives more information about people responding [BBBB] than can 3-PL.

## General Discussion

The model proposed here provides the first practical, complete IRT item analysis for multiple choice tests, describing the performance of all of the response alternatives as functions of the trait being measured. The parameters of the model are not readily interpretable for the most part, but graphical presentation of the trace lines and/or their products with the population distribution gives thorough item analysis which has much to recommend it.

This sort of item analysis is sufficiently flexible that "bad" trace lines are fitted. These can then be observed and such items modified or eliminated in the course of empirical test development. That is not possible, in general, with simpler models which require examination of the residuals to find bad items. For item analysis, the model of (2) and (5) is clearly an excellent choice; for scoring tests (aka "estimating  $\hat{\theta}$ "), matters are more complicated.

Non-monotonic trace lines for the correct response have become a popular feature of recent IRT developments: Lord (1982) and Choppin

(1983) have proposed item response models which permit "low-ability" non-monotonicity, in work on binary models independent from Samejima's (1979) multiple category proposal. There are a number of possible explanations for non-monotonicity on the left, as for the correct response in Figure 1. The only requirement for candidacy as an explanation for the effect is that it must account for "getting the right answer for the wrong reason." Bock (personal communication, 1983) suggests a name for the phenomenon: "positive misinformation."

Two sources of positive misinformation come to mind. The first is that the correct response for a particular item differs from the distractors on dimensions other than that which is intended and observable, given sufficient ability. Examinees of medium and high ability perceive the features and attempt the processing intended by the item-writer, while examinees of low ability see other features of the alternatives which cause them to select the correct one. These "other features" may be effectively "invisible" to individuals of higher ability, and therefore to the item-writers as well.

A second possible cause of non-monotonicity in the correct trace line is cheating. If low-ability individuals cheat (e.g., by copying a neighbor's answer, which is more likely to be correct than not for most items), then the resulting correct alternative trace lines will rise for those of very low ability.

It may also be possible that non-monotonic trace lines are estimation artifacts in small sets of items (like four). It may be that, with few monotonic correct-alternative trace lines to "orient"  $\theta$ , the estimative trace lines trace

tion procedure could "wrap around" and (effectively) place some of the high ability individuals on the left end of the  $\theta$ -continuum. But this explanation can be discounted with longer tests, as the non-monotonicity there comes from correct response to an item being more likely from examinees who get most of the other items wrong than from those who get fewer of the other items wrong.

Distinguishing among these possibilities would require experimentation with the items and the testing situation. The model does, however, offer an item analysis which permits such phenomena to exhibit themselves in the fit, when they are present.

So the use of non-monotonic curves in item analysis may be required to fit the observed data. However, the use of non-monotonic trace lines in constructing the posterior density, a measure of the central tendency of which is to be called  $\hat{\theta}$  and used to "score the test", gives rise to many potential problems. If the correct response trace line for item j is non-monotonic on the right (e.g., it turns down), then examinees with some response patterns on the other items will be "penalized" by responding correctly to item j; they would have been assigned a higher  $\hat{\theta}$  if they had selected certain of the incorrect alternatives. Correct response trace lines which are non-monotonic on the left similarly "penalize" examinees of low ability who respond correctly. As measurement, this is all probably satisfactory: conditional on the other item responses, a correct response to item j may not imply higher ability; it may be more likely to be guessing or cheating. But this may be a problem for the test "as contest." Further, it might be difficult

to defend such a method of test scoring against the onslaught of members of the bar before a jury. A case can, therefore, be made for either a) rejecting the model or b) rejecting the use of items with non-monotonic trace lines for the correct response.

Multiple category scoring of this nonlinear sort has certain bizarre properties even if non-monotonic trace lines are eliminated. For certain regions on the  $\theta$ -continuum, selecting a particular incorrect response will increase  $\hat{\theta}$  more than would selecting the actual correct response. For instance, in the Word Knowledge item in Figure 2, for  $\theta$  just below -1, selecting response B will increase  $\hat{\theta}$  more than would selecting the correct response. But for  $\theta$  around 1, selecting B will "penalize" the respondent! The contingencies are sufficiently complex that it is unlikely that the examinees could find a strategy to take advantage of the system. But the possibilities for legal difficulties are considerable under circumstances in which the test and scoring system must be disclosed. On the other hand, the quality of measurement for research purposes and in non-disclosed tests should be improved by these "bizarre" features; that is, after all, where the multiple category model obtains its additional information.

## Conclusion

Item response models for multiple choice items have come of age: they fit the data. Questions remain about uses for these models. They produce excellent item analysis, but it is complex and best-represented graphically -- that breaks with the tradition of item analysis with a

few numerical summaries. Multiple category scoring clearly increases the information obtained in scoring a test -- but again at the cost of complexity, and, potentially, controversy. The validity of test scored with such methods has not been examined here, nor can it be with internal consistency data in any event. Consideration of all of these matters is deferred to future work.

#### References

Bock, R.D. Estimating item parameters and latent ability when the responses are scored in two or more nominal categories. <u>Psychometrika</u>, 1972, 37, 29-51.

Bock, R.D. and Aitkin, M. Marginal maximum likelihood estimation of item parameters: An application of an EM algorithm. <u>Psychometrika</u>, 1981, 46, 443-459.

Bock, R.D. and Lieberman, M. Fitting a response model for n dichotomously scored items. Psychometrika, 1970, 35, 179-197.

Bock, R.D. and Mislevy, R.G. Data quality analysis of the Armed Forces Vocational Aptitude Battery. DOD MDA 903-802-0333. National Opinion Research Center, Chicago, 1981.

Bock, R.D. and Mislevy, R.G. Applications of EAP estimation in computerized adaptive testing. Paper presented at the annual meeting of the Psychometric Society, Montreal, 1982.

Choppin, B. A two-parameter latent trait model. CSE Report no. 197, Center for the Study of Evaluation, Graduate School of Education, University of California, Los Angeles, 1983.

LESSESSES TRANSPORTED TO SERVICE SERVICE TO SERVICE SE

Dempster, A.P., Laird, N.M., and Rubin, D.B. Maximum likelihood from incomplete data via the EM algorithm (with Discussion). <u>Journal of the Royal Statistical Society</u>, Series B, 1977, 39, 1-38.

Haberman, S. Subroutine MINIM. In R. D. Bock and Bruno Repp (Eds.) MAT-CAL: Double precision matrix operations subroutines for the IBM System 360/370 computers. Chicago: National Educational Resources, 1974.

Levine, M.V. and Drasgow, F. The relation between incorrect option choice and estimated ability. Educational and Psychological Measurement, 1982, in press.

Lord, F.M. Maximum likelihood estimation of item response parameters when some responses are omitted. In preparation, 1982.

Samejima, F. A new family of models for the multiple choice item.

Research Report #79-4, Department of Psychology, University of Tennessee, 1979.

Sympson, J.B. A new IRT model for calibrating multiple choice items.

Paper presented at the annual meeting of the Psychometric Society, Los

Angeles, 1983.

Thissen, D. Information in wrong responses to the Raven Progressive Matrices. <u>Journal of Educational Measurement</u>, 1976, 13, 201-214.

Thissen, D. Marginal maximum likelihood estimation for the one-parameter logistic model. <u>Psychometrika</u>, 1982, 47, 175-186.

## Footnotes

This research was supported in part by ONR contract N00014-83-C0283 to the University of Chicago, R. Darrell Bock, Principal Investigator
and AFHRL contract F41689-82-C-00020 to the Educational Testing Service,
Howard Wainer, Principal Investigator. This project was facilitated by
a sabbatical leave provided DT by the University of Kansas, during which time
some of this work was done with the Research Statistics Group of the
Educational Testing Service. Conversations with Darrell Bock, Howard
Wainer, Paul Holland, Frederic Lord, Donald Rubin and James Ramsay have
also been important and useful in shaping the research and its presentation
here. While all of those mentioned have helped us gain wisdom in these
matters, any error that remains is, of course, our own.

Table 1.

Cross classification of responses to two items (22 and 23)

of the ASVAB Form 8A General Science subtest for 1048 examinees

of the NORC national probability sample (Bock and Mislevy, 1981).

		item 2				
		A	В	C؉	D	
	A	20(22)	107(23)	79(22)	44(31)	250
Item	B* .	32(36)	157(34)	143(40)	35(25)	367
1	С	7(8)	36(8)	24(7)	19(14)	86
	D	30(33)	161(35)	112(31)	42(30)	345
		89	461	358	140	1048

\*=Correct;

Column percentages in parentheses.

Table 2.  $\label{eq:G2} \textbf{G}^{\textbf{2}} \ \, \textbf{values} \ \, \textbf{for selected models for the}$   $\label{eq:G2} \quad \, \textbf{four-item examples}.$ 

		Word		General	
		Knowledge		Science	
Model	d.f.	G <sup>2</sup>	p	G²	p
Bock (1972)	231	307	<.01	277	<.02
Samejima (1979)	223	271	<.02	264	<.05
"ABCD"	220	258	<.04	262	<.02
"ABCD(C),ABCD(D)"	217	248	>.05		
"11 per item"	211	245	<.05		
"Correct vs. incorrect"	222			253	≃.05
"Correct vs., each item"	3			253	= 05

Table 3.

Estimated parameters for the last four Word Knowledge items.

Estimates for four items above, whole test below.

Parameters for correct response underscored.

		Response				
		DK	A	В	C	D
Item 32	a k	-1.1	-2.9	1.5	0.2	2.2
	c <sub>k</sub>	-1.7 0.1 0.3	-1.0 -3.8 -2.3 0.1 0.25	1.1 2.9 2.4 0.2	0.3 -1.9 -2.5 0.3	$\frac{1.9}{2.7}$ $\frac{2.1}{0.4}$
33	ak ck dk	-3.0 -2.1 0.1 2.1	-0.3 -0.6 0.6 0.5 0.2	0.25 1.2 1.2 -1.6 -3.0 0.2 0.2	$ \begin{array}{r} 0.25 \\ \underline{2.8} \\ \underline{2.3} \\ 0.1 \\ -0.6 \\ \underline{0.5} \\ 0.4 \end{array} $	0.25 -0.7 -0.8 0.8 1.0 0.1
34	ak ck dk	-0.8 -1.3 -1.6 -0.9	-1.8 -0.9 -3.2 -2.5 0.2	-0.1 -0.2 0.2 -0.1 0.2	$   \begin{array}{r}     1.9 \\     \hline     1.9 \\     \hline     2.6 \\     \hline     1.8 \\     \hline     0.5 \\     0.4 \\   \end{array} $	0.2 0.7 0.5 2.0 1.6 0.1 0.2
35	ak ck dk	-1.6 -1.9 0.0 -0.1	-0.4 0.5 -1.3 -2.0 0.1 0.25	-0.2 0.0 0.4 0.5 0.2 0.25	-0.8 -0.6 0.4 0.8 0.3 0.25	3.0 1.9 0.5 0.8 0.4 0.25

Table 4.

Estimated parameters for the last four General Science items.

Estimates for four items above, "physical science" subset below.

Parameters for correct response underscored.

		Response				
		DK	A	В	С	D
Item 22	a k	-2.3	-0.2	2.0	0.9	-0.3
	c k	0.5	0.7	- <u>0.5</u>	-1.9	1.1
	<sup>d</sup> k		0.2	0.4	0.2	0.2
23	ak ck dk	-0.8 -4.0 1.6 -0.8	0.6 1.2 -2.8 -0.7 0.2 0.2	-0.5 1.1 1.5 1.4 0.2 0.3	$\begin{array}{r} 1.1 \\ 2.5 \\ \hline 0.0 \\ \hline 0.2 \\ \hline 0.4 \\ \hline 0.3 \end{array}$	-0.4 -0.9 -0.3 -0.1 0.2 0.2
24	ak ck dk	-0.5 -3.3 -0.3 -1.8	-0.2 0.2 0.7 1.1 0.2 0.2	$\begin{array}{r} 2.0 \\ \hline 2.6 \\ -\overline{1.0} \\ -\overline{1.0} \\ \hline 0.4 \\ \hline 0.4 \end{array}$	-1.2 -0.1 0.7 1.4 0.2 0.2	0.0 0.5 0.0 0.2 0.2
25	a <sub>k</sub>	-1.5	-0.7	-0.2	0.1	2.3
	ck	0.4	0.4	-0.5	0.5	<u>-0.8</u>
	$\mathbf{d}_{\mathbf{k}}$		0.2	0.2	0.2	0.4

## Figure Captions


Figure 1. Five panels showing P(k),  $k=0,1,\ldots,m_j$  as a function of  $\theta$ ; the solid lines correspond to the probabilities for DK, and responses A,B,C, and D for GS item 23, estimated as part of "physica' science." The dashed curves use the estimates obtained with the last four items.


Figure 2. Five panels showing P(k),  $k=0,1,\ldots,m_j$  as a function of  $\theta$ ; the solid lines correspond to the probabilities for DK, and responses A,B,C, and D for WK item 32 estimated in the entire test. The dashed curves use the estimates obtained with the last four items.


Figure 3. Four panels show three alternative fits of the trace line for the correct responses for the last four WK items: dots are on the standard 3-PL curve, the solid line is the model of the present paper fitted with the entire test, and the dashed line is the present model fitted with only those items.


Figure 4. Test information curves for binary (dotted line) and multiple category (solid line) scoring for WK, as functions of  $\theta$ .


Figure 5. The solid lines give the trace lines from the present model and posterior density (labelled "Total") for response pattern [BBBB] (all incorrect) for the last four items of WK. Dashed lines give the corresponding curves for the same response pattern using 3-PL estimates.











Lowell Schoer
Psycho. & Quanti. Foundations
College of Education
University of lowa
lowa City, IA 52242

Dr. Edwin Shirkey
Department of Psychology
Univer. of Central Florida
Orlando, FL 32816

Dr. William Sims Center for Naval Analysis 200 North Beauregard Street Alexandria, VA 22311

Dr. H. Wallace Sinaiko Smithsonian Institution 801 North Pitt Street Alexandria, VA 22314

Dr. Kathryn T. Spoehr Psychology Department Brown University Providence, RI 02912

Dr. Robert Sternberg
Department of Psychology
Yale University
Box 11A, Yale Station
New Haven, CT 06520

Dr. Peter Stoloff Cenzer for Naval Analysis 200 North Beauregard Street Alexandria, VA 22311

Dr. William Stout University of Illinois Department of Mathematics Urbana, IL 61801

Dr. Patrick Suppes
Insti. for Mathematical Studies
In the Social Sciences
Stanford University
Stanford, CA 94305
Dr. Kurt Va
Xerox PARC
3333 Coyota
Palo Alto,

Dr. Hariharan Swaminathan Lab. of Psy. & Eval. Research School of Education University of Massachusetts Amberst, MA 01003 Dr. Kikumi Tatsuoka Comp. Based Educ. Research 252 Eng. Research Laboratories Urbana. IL 61820

Dr. Maurice Tatsuoka 220 Education Building 1310 South Sixth Street Champaign, 1L 61820

Dr. David Thissen
Department of Psychology
University of Kansas
Lawrence, KS 66044

Or. Douglas Towne Univer. of So. California Behavioral Technology Labs 1845 S. Elena Avenue Redondo Beach, CA 90277

Dr. Robert Tsutakawa Department of Statistics University of Missouri Columbia, MO 65201

Or. J. Uhlaner Uhlaner Consultants 4258 Bonavita Orive Encino, CA 91436

Or. V. R. R. Uppuluri Union Carbide Corporation Nuclear Division P.O. Box Y Oak Ridge, TN 37830

Dr. David Vale
Assessment Systems Corporation
2233 University Avenue
Suite 310
St. Paul, MN 55114

Dr. Kurt Van Lehn Xerox PARC 3333 Coyote Hill Road Palo Alto, CA 94304

Dr. Howard Wainer
Division of Psy. Studies
Educational Testing Service
Princeton, NJ 08540

Or. Michael T. WAller Department of Educ. Psychology University of Wis.-Milwaukee Milwaukee, WI 53201

Dr. Brian Waters HumRRO 300 North Washington Alexandria, VA 22314

Or. Phyllis Weaver 2979 Alexis Orive Palo Alto, CA 94304

Dr. David J. Weiss N660 Elliott Hall University of Minnesota 75 E. River Road Minneapolis, MN 55455

Dr. Donald O. Weitzman Mitre Corporation 1820 Dolley Madison Blvd McLean, VA 22102

Or. Rand R. Wilcox University of Southern Califor Department of Psychology Los Angeles, CA 90007

Wolfgang Wildgrube Streitkraefteamt Box 20 50 03 D-5300 Bonn 2 WEST GERMANY

Dr. Bruce Williams Depart. of Educ. Psychology University of Illinois Urbana, IL 61801

Or. Wendy Yen CTB/McGraw Hill Del Monte Research Park Monterey, CA 93940

H. William Greenup Education Advisor (E031 Education Center MCDEC Quantico, VA 22134

Or. Douglas H. Jones Advanced Statistical Technologies Corporation 10 Trafalgar Court Lawrenceville, NJ 08148

Professor John A. Keats Department of Psychology The University of Newcastle N.S.W. 2308 **AUSTRALIA** 

Dr. Marcy Lansman University of North Carolina Davie Hall 013A Capel Hill, NC 27514

Dr. Alan Lesgold Learning R&D Center University of Pittsburgh 3939 O'Hara Street Pittsburgh, PA 15260

edition therefore the common the common the parameter of

Dr. Michael Levine Depart. of Educa. Psychology 210 Education Bldg. University of Illinois Champaign, IL 61801

Dr. Charles Lewis Fac. Soc. Wetenschappe Rijksuniversiteit Groningen Oude Boteringestraat 23 3712GC Groningen, NETHERLANDS

Dr. Robert Linn College of Education University of Illinois Urbana, IL 61801

Mr. Phillip Livingston Systems & Applied Sci. Corp. 6811 Kenilworth Avenue Riverdale, MD 20840

Dr. Robert Lockman Center for Naval Analysis 200 North Beauregard Street Alaexandria, VA 22311

Dr. Frederic M. Lord Educational Testing Service Princeton, NJ 08541

Dr. James Lumsden Department of Psychology University of Western Australia Next ands W.A. 6009 AUSTRALIA

Or. Gary Marco Stop 31-E Educational Testing Service Princeton, NJ 08451

the Contract of the Contract o

Dr. Scott Maxwell Department of Psychology University of Notre Dame Notre Dame, IN 46556

Or. Samuel T. Mayo Loyola University of Chicago 820 North Michigan Avenue Chicago, IL 60611

Mr. Robert McKinley American College Testing Pro. P.O. Box 168 lowa City, IA 52243

Dr. Barbara Means Human Resources Research Organ. 300 North Washington Alexandria, VA 22314

Dr. Allen Munro Behavioral Tech. Laboratories 1845 Elena Ave., Fourth Floor Redondo Beach, CA 90277

Dr. W. Alan Nicewander University of Oklahoma Department of Psychology Oklahoma City, OK 73069

Dr. Donald A. Norman Cognitive Science, C-015 Univ. of California, San Diego La Jolla, CA 92093

Dr. Melvin R. Novick 356 Lindquist Cnt. for Measure. Department of Psychology University of lowa lowa City, IA 52242

Dr. James Olson WICAT, Inc. 1875 Wouth State Street Orem, UT 84057

Wayne M. Patience American Council on Education GED Testing Service, Suite 20 One Dupon Cirle, NW Washington, DC 20036

Dr. James A. Paulson Portland State University P.O. Box 751 Portland, OR 97207

Dr. James W. Pellegrino University of California, Santa Barbara Department of Psychology Santa Barabara, CA 93106

Dr. Mark D. Reckase P.O. Box 168 lowa City, IA 52243

Dr. Lauren Resnick LRDC University of Pittsburgh 3939 O'Hara Street Pittsburg, PA 15260

Dr. Thomas Reynolds University of Texas-Dallas Marketing Department P.O. Box 688 Richardson, TX 75080

Dr. Andrew M. Rose Amer. Insti. for Research 1055 Thomas Jefferson St. NW Washington, DC 20007

Dr. Lawrence Rudner 403 Elm Avenue Takoma Park, MD 20012

Dr. J. Ryan Department of Education University of South Carolina Columbia, SC 29208

Prof. Fumiko Samejima Univeristy of Tennessee Knoxville, TN 37916

Dr. Walter Schneider Psychology Department 603 E. Daniel Champaign, 1L 61820

Or. Kazuo Shigemasu 7-9-24 Kugenuma-Kaigan Fujusawa 251 JAPAN

Dr. Richard L. Ferguson Dr. Allan M. Collins Dr. Erling B. Andersen The American College Testing Boit Beranek & Newman, Inc. Department of Statistics . Program 50 Moulton Street. Studiestraede 6 P.O. Box 168 Cambridge, MA 02138 1455 Copenhagen lowa City, 1A 52240 DEMMARK Univ. Prof. Dr. Gerhard Dr. Isaac Bejar Dr. Hans Crombag Fischer Educational Testing Service Education Research Center Liebiggasse 5/3 Princeton, NJ 08450 University of Leyden A 1010 Vienna Boerhaavelaan 2 AUSTRIA 2334 EN Leyden, NETHERLANDS Professor Donald Fitzgerald CTB/McGraw-Hill Library Dr. Menucha Birenbaum University of New England 2500 Garden Road School of Education Armidale, New South Wales Monterey, CA 93940 Tell Aviv University 2351 Tell Aviv, Ramt Aviv 60078 AUSTRALIA ISRAEL Dr. Dexter Fletcher Dr. Werner Birke Dr. Dattpradåd Divgi University of Oregon Personalstammamt der Bundeswehr Syracuse University Department of Computer Science Department of Psychology D-5000 Hoein 90 Eugene, OR 97403 Syracuse, NE 33210 WEST GERMANY · Dr. John R. Rfederiksen Dr. Robert Brennan Dr. Fritz Drasgow Bolt Beranek & Newman American College Testing Pro-Department of Psychology 50 Moulton Street Univeristy of Illinois crams Cambridge, MA 02138 P.O. Box 168 603 E. Daniel St. Lowa City, IA 52243 Champaign, IL 61820 Dr. Susan Embertson Dr. Janice Gifford Dr. Glenn Bryan University of Massachusetts Psychology Department 6208 Poe Road University of Kansas School of Education Bethesda, MD 20817 Lawrence, KS 66045 Amherst, MA 01002 ERIC Facility-Acquisitions Dr. Robert Glaser Dr. Ernest R. Cadotte 4833 Rugby Avenue Learning Research & Dev. Cente 307 Stokely University of Tennnessee Bethesda, MD 20014 University of Pittsburgh 3939 O'Hara Street Knoxville, TN 37916 Pittsburgh, PA 15260 Or. Benjamin A. Fairbank, Jr. Dr. Bert Green Dr. John B. Carroll McFann-Gray & Associates, Inc. Johns Hopkins University 409 Elliott Road 5825 Callaghan Department of Psychology Chapel Hill, NC 27514 Suite 225 Charles & 34th Street San Antonio, TX 78228 Baltimore, MD 21218 Dr. Ron Hambleton Dr. Leonard Feldt Dr. Norman Cliff School of Education Lindquist Center for Measure-Department of Psychology University of Massachusetts ment Univ. of So. California Amherst, MA 01002 Univesity of lowa Univesity Park lowa City, IA 52242 Los Angeles, CA 90007 Dr. Paul Horst Dr. Lloyd Humphreys Dr. Earl Hunt 677 G Street, #184 Department of Psychology Department of Psychology Chula Vista, CA 90010 University of Washington University of Illinois 603 East Daniel Street Seattle, WA 98105 Champaign, IL 61820 Dr. William Koch Dr. Huynh Huynh Dr. Jack Hunter University of Texas-Austin 2122 Coolidge Street College of Education

Lansing, MI 48906

University of South Carolina

Columbia, SC 29208

Measurement and Evaluation

Austin, TX 78703

Center

ATR FORCE HUMAN RESOURCES LAB AFHRL/MPO

Technical Documents Center US AF OFFICE OF SCI. RES.

AF Human Resources Lab LIFE SCI. DIRECTORATE, NL
WPAFB, OH 45433 BOLLING AIR FORCE BASE AFHRL/MPO AF Human Resources Lab BROOKS AFB, TX 78235 WPAFB, OH 45433

US AF OFFICE OF SCI. RES. WASHINGTON, DC 20332

AIR UNIVERSITY LIBRARY AUL/LSE 76/443 MAXWELL AFB, AL 36112

OR. EARL A. ALLUISI MR. RAYMOND E. CHRISTAL HQ, AFHRL (AFSC) AFHRL/MOE BROOKS AFB, TX 78235 BROOKS AFB, TX 78235

DR. ALFRED R. FREGLY AFOSR/NL BOLLING AFB, DC 20332

DR. GENEVIEVE HADDAD DR. T. M. LONGRIDGE
PROGRAM MANAGER AFHRL/OTE
LIFE SCIENCES DIRECTORATE WILLIAMS AFB, AZ 85224
AFOSR BOLLING AFB. DC 20332

MR. RANDOLPH PARK DR. ROGER PENNELL DR. MALCOM REE
AFHRL/MOAN AF HUMAN RESOURCES LAB
BROOKS AFB, TX 78235 LOWRY AFB, CO 80230 BROOKS AFB, TX 78235

3700 TCHTW/TTGHR

2LT TALLARIGO

SHEPPARD AFB, TX 76311

LT. COL JAMES E. WATSON

HQ USAF/MPXOA

THE PENTAGON

BROOKS AFB, TX WASHINGTON, DC 20330

DR. JOSEPH YASATUKE AFHRL/LRT LOWRY AFB, CO 80230

DEFENSE TECHNICAL INFO CEN.

CAMERON STATION, BLDG 5

ALEXANDRIA, VA 22314

ATTN: TC

DR. CRAIG I. FIELDS

ADVANCED RES. PROJECTS AGENCY

1400 WILSON BLVD.

ARLINGTON, VA 22209

DR. WILLIAM GRAHAM TESTING DIRECTORATE MEPCOM/MEPCT-P FT. SHERIDAN, IL 60037

Jerry Lehnus HQ MEPCOM Arth ....

Jerry Lehnus MILITARY ASS. FOR TRAINING 5
HQ MEPCOM PERSONNEL TECH.
Attn: MEPCT-P OFF. OF THE UNDER SEC. FOR REFORM 30129, the PENTAGON WASHINGTON, DC 20301 WASHINGTON, DC 20301

DR. WAYNE SELLMAN

OFF. OF THE ASS. SEC. OF

DEFENSE (MRA ε L)

28269 the Pentagon

ARLINGTON, VA 22209

DR. SUSAN CHIPMAN

LEARNING ε DEVELOPMENT

NAT'L INST. OF EDUCATION

1200 19th STREET NO WASHINGTON, DC 20301

WASHINGTON, DC 20208

DR. ARTHUR MELMED

DR. ANDREW R. MOLNAR

724 BROWN

US DEPARTMENT OF ED.

WASHINGTON, DC 20208

DR. ANDREW R. MOLNAR

OFF. OF SCI & ENG PER & ED

PERSONNEL R&D CENTER

OFF. OF PERSONNEL MANAGEMENT

WASHINGTON, DC 20208

WASHINGTON, DC 20415

MR. THOMAS A. WARM
US COAST GUARD INST.
P.O. SUBSTATION 18
DKLAHOMA CITY, OK 73169

DR. JOSEPH L. YOUNG, DIR.
MEMORY & COGNITIVE PROCESSES
UNIVERSITY OF FLOW
GAINESVILLE, FL 33
WASHINGTON, DC 20550

UNIVERSITY OF FLORIDA GAINESVILLE, FL 326

PERSONNEL & TRAINING RESEARCH GROUP ARLINGTON, VA 22217

PENSACOLA, FL 32508 WASHINGTON, DC 20350

DR. CARL ROSS
CNET-PDCD BUILDING 90 GREAT LAKES NTC, IL 60088

DR. R. SNOW, LIAISON SCI.

OFFICE OF NAVAL RESEARCH
BRANCH OFFICE, LONDON

DR. RICHARD SORENSEN

NAVY PERSONNEL RED CENTER - CNO - OP115

SAN DIEGO, CA 92152

NAVY ANNEX

ARLINGTON, VA BOX 39 FPO NEW YORK, NY 09510

UR. DOUGLAS WETZEL

CODE 12

NAVY PERSONNEL RED CENTER
SAN DIEGO, CA 92152

OR. MARTIN F. WISKOFF
NAVY PERSONNEL RED CENTER
SAN DIEGO, CA 92152 DR. DOUGLAS WETZEL

DR. ROBERT CARROLL NAVOP 115 WASHINGTON, DC 20370

DEPT. OF ADM. SCIENCES CODE P13
NAVAL POSTGRADUATE SCHOOL NPRDC
MONTEREY, CA 93940

MR. BRAD SYMPSON

OR. MARTIN A. TOLCOTT

NAVY PERSONNEL R&D CENTER

SAN DIEGO, CA 92152

OR. JAMES TWEEDDALE

TECHNICAL DIRECTOR

NAVY PERSONNEL R&D CENTER

SAN DIEGO, CA 92152

ARLINGTON, VA 22217

SAN DIEGO, CA 92152

RESEARCH GROUP

CODE 442PT

OFFICE OF NAVAL RESEARCH

AND TRAINING (OP-01E)

RM. 2705 ARLINGTON ANNEX

WASHINGTON, DC 20370

LT FRANK C. PETHO, MSC,

USN (PH.D)

CNET (N-432)

MAS

OFFICE OF THE CHIEF OF NAVAL

OPERATIONS

RES. DEV. & STUDIES BRANCH

OP 115

DR. FRANK VICINO

NAVY PERSONNEL RED CENTER

SAN DIEGO, CA 92152

OR. EDWARD WEGMAN

OFFICE OF NAVAL RESEARCH

SOU NORTH QUINCY STREET

ARLINGTON, VA 22217

DR. RONALD WEITZMAN

NAVAL POSTGRADUATE SCHOOL

DEPT. OF ADM. SCIENCES

MONTEREY, CA 93940

DR. WALLACE WULFECK, III

DR. NICK BOND

NAVY PERSONNEL R&D CENTER

SAN DIEGO, CA 92152

DR. NICK BOND

OFFICE OF NAVAL RESEARCH

LIAISON OFFICE, FAR EAST

MEASUREMENT DIVISIO APO SAN FRANCISCO, CA 96503

> DR. ROBERT BREAUX NAVTRAEQUIPCEN CODE N-095R ORLANDO, FL 32813

> > ARLINGTON, VA 22217

SPECIAL ASST. FOR EDUCATION GEREXERXMEXIMEXEMISEXCEXMENTAL OPERATIONS-RESEARCH-BEYELOPMENT-5 ---STUB+ES-BRANEH-

> DR. BERNARD RIMLAND (01C) NAVY PERSONNEL RED CENTER SAN DIEGO, CA 95152

OR. ROBERT G. SMITH

OFFICE OF CHIEF OF NAVAL OP.

OP-987H

WASHINGTON, DC 20350

DR. ALFRED F. SMODE, DIR.

TRAINING ANALYSIS &

EVALUATION GROUP

DEPT. OF THE NAVY

ORLANDO, FL 32813

DR. FREDERICK STEINHEISER ARLINGTON, VA 20370

MR. JOHN H. WOLFE NAVY PERSONNEL RED CENTER SAN DIEGO, CA 92152

MEASUREMENT DIVISION NAS PENSACOLA, FL 32508

Chief of Naval Ed. & Training Liason Office Air Force Human Resource Lab Operations Training Division WILLIAMS AFB, AZ 85224

DR. STANLEY COLLYER

OFFICE OF NAVAL TECHNOLOGY

800 N. QUINCY ST.

ARLINGTON, VA 22217

CDR MIKE CURRAN

PROGRAM DEVELOPMENT

SUILDING 90

NET-PDCD

ARLINGTON, VA 22217

DR. CATHY FERNANDES NAVY PERSONNEL RED CENTER SAN DIEGO, CA 92152

MR.	PAUL	FOLEY		
MAVY	PERS	ONNEL	R&D	CENTER
SAN	01500	, SA	9215	52

DR. WILLIAM L. MALOY (20)

CMIEF OF NAVAL ED. 8

TRAINING

TRAINING

SAN DIESO, CA 92152

DR. WILLIAM HONTAGUE
NPRDC CODE 13
SAN DIESO, CA 92152

SAN DIESO, CA 92152 PENSACOLA, FL 32508

DR. WILLIAM E. NORDBROCK Library. Code P201L THC-ADCO BOX 25 APO NY 09710

STEC. ASS. FOR MARINE MATTERS OR. A.L. SLAFKOSKY CODE 1004 OFFICE OF NAVAL RESEARCH

BOD N. QUINCY STREET

ABILINETON VA 22217

ABILINETON VA 22217

CODE MP1-20)

WASHINGTON, DC 20380

WASHINGTON, DC 20380

TECHNICAL DIRECTOR

U.S. ARMY RESEARCH INST.

BEHAVORIAL S SOCIAL SCIENCES
5701 EISENHOWER AVENUE
ALEXANDRIA, VA 22333

MR. JAMES BAKER

DR. KENT BATON

ARMY RESEARCH INSTITUTE
SOOI EISENHOWER AVENUE
ALEXANDRIA, VA 22333

ALEXANDRIA, VA 22333

MR. ROBERT ROSS THE ROBERT ROSS

DR. ROBERT SASNOR

DR. JOYCE SHIELDS

LE ARMY RESEARCH INST. FOR THE US GENY RES. INST. FOR THE STOCIAL & SEMAVIORAL & SOCIAL SCIENCES

STOT EISENHOWER AVENUE SOCIAL SCIENCES

ALEXANDRIA, VA 22333

DR. ROBERT SASNOR

DR. JOYCE SHIELDS

ARMY RESEARCH INST. FOR THE SEMAVIORAL & SOCIAL SCIENCES

SCHAVIORAL & SOCIAL SCIENCES

SOCIAL SCIENCES

ALEXANDRIA, VA 22333

ALEXANDRIA, VA 22333

DR. ROBERT J. SEIDEL US ARMY RESEARCH INST. 5001 EISENHOWER AVINUE 3000 N. WASHINGTON ST. ALEXANDRIA, VA 22333

dr. jim hollan

DR. NORMAN J. KERR DR. LEONARD KROEKER
MAVY PERSONNEL RED CENTER CHIEF OF NAVAL TECH. TRAINING NAVY PERSONNEL RED CENTER
SAN DIEGO, CA 92132 NAVAL AIR STATION MEMPHIS (75) SAN DIEGO. CA 92152 MILLINGTON, TN 38054

Library. Code P201L TECHNICAL DIRECTOR
Navy Personnel R&D Center MAVY PERSONNEL R&D CENTER
San Diego. CA 92152 SAM DIEGO, CA 92152

H\_ WILLIA'S GREENUP

EDUCATION ADVISOR (E031)

EDUCATION CENTER, MCDEC

QUANTICO, VA 22134

DIR., DFFICE OF MANPOWER

UTILIZATION

HQ. MARINE CORPS (MPU)

ECS. ELDG. 2009 QUANTICO, VA 22134

SCIENTIFIC ADVISOR (CODE RD-1) HEADQUARTERS. MARINE CORPS

DR. BEATRICE J. FARR
U.S. ARMY RESEARCH INST.
U.S. ARMY RESEARCH INST.

5001 EISENHOWER AVENUE
ALEXANDRIA, VA 22333

DR. MYRON FISCHL
U.S. ARMY RESEARCH INST.

TRAINING TECHNICAL AREA
U.S. ARMY RESEARCH INST.

5001 EISENHOWER AVENUE
5001 EISENHOWER AVENUE
ALEXANDRIA, VA 22333

ALEXANDRIA, VA 22333

DR. HARDLD F. C'NEIL, JR.

DIRECTOR. TRAINING RES. LAB

FOR BEHAVIORAL & SOCIAL SCI.

ATTN: PERI-IC

ACTN: PERI

DR. ROBERT SASMOR

DR. HILDA WING ARMY RESEARCH INST. **5001 Eisenhower Ave**nue Alexandria, va 22333

DR. JIH HOLLAN COUC 14 HAVY PERSONNEL ROD CENTER SAN DIEGO, CA 92152

TECHNICAL DIRECTOR

Headquarters, US Marine Corps 🔄 Code MP1-20 Washington, DC 20380

MAJOR FRANK YOHANNAN. USMC

OR. JOYCE SHIELDS

DR. ROBERT WISHER ARMY RESEARCH INST. 5001 EISENHOWER AVE. ALEXANDRIA, VA 22333

FILMED

ACA

DIFFIC